

Systematic Comparison of Jet Energy-Loss Schemes in a 3D hydrodynamic medium

S.A. Bass¹, C. Gale², A. Majumder¹, C. Nonaka³, G.-Y. Qin²,
T. Renk^{4,5} and J. Ruppert^{2,6}

¹ Department of Physics, Duke University, Durham, North Carolina 27708-0305, USA

² Department of Physics, McGill University, H3A 2T8, Montreal, Quebec, Canada

³ Department of Physics, Nagoya University, Nagoya 464-8602, Japan

⁴ Department of Physics, PO Box 35 FIN-40014 University of Jyväskylä, Finland

⁵ Helsinki Institute of Physics, PO Box 64 FIN-00014 University of Helsinki, Finland

⁶ Institut für Theoretische Physik,, J.W. Goethe University Frankfurt,
Max-von-Laue-Str. 1, D-60438 Frankfurt am Main, Germany

E-mail: bass@phy.duke.edu

Submitted to: *J. Phys. G: Nucl. Phys.*

PACS numbers: 25.75.-q, 12.38.Mh

Abstract. We utilize a 3D hydrodynamic model to provide the medium evolution for a systematic comparison of jet energy-loss calculations in the BDMPS/ASW, HT and AMY approaches. We find that the parameters of all three calculations can be adjusted to provide a good description of inclusive data on R_{AA} versus transverse momentum. However, we do observe slight differences in their predictions for the azimuthal angular dependence of R_{AA} vs. p_T . We also note that the value of the transport coefficient $\hat{q}$ needed in the three approaches to describe the data differs significantly.

Experiments at the Relativistic Heavy Ion Collider (RHIC) have established a significant suppression of high- p_T hadrons produced in central A+A collisions compared to those produced in peripheral A+A or binary scaled p+p reactions, indicating a strong nuclear medium effect [1, 2], commonly referred to as *jet-quenching*. Within the framework of perturbative QCD, the leading process of energy loss of a fast parton is gluon radiation induced by multiple soft collisions of the leading parton or the radiated gluon with color charges in the quasi-thermal medium [3, 4, 5].

Over the past two years, a large amount of jet-quenching related experimental data has become available, including but not limited to the nuclear modification factor R_{AA} , the elliptic flow v_2 at high p_T (as a measure of the azimuthal anisotropy of the jet cross section) and a whole array of high p_T hadron-hadron correlations. Computations of such jet modifications have acquired a certain level of sophistication regarding the incorporation of the partonic processes involved. However, most of these calculations have been utilizing simplified models for the underlying soft medium, e.g. assuming




Figure 1. Left: Nuclear modification factor R_{AA} in $Au-Au$ collisions at 0-5% (top) and 20-30% (bottom) centrality calculated in the ASW, HT and AMY approaches compared to data from PHENIX [12]. Right: R_{AA} as a function of azimuthal angle at $p_T = 10$ GeV/c (solid line) and $p_T = 15$ GeV/c (dashed line) for all three approaches in the 20-30% centrality bin.

a simple density distribution and its variation with time. Even in more elaborate setups, most jet quenching calculations assume merely a one- or two-dimensional Bjorken expansion.

The availability of a three-dimensional hydrodynamic evolution code [6] allow for a much more detailed study of jet interactions in a longitudinally and transversely expanding medium. The variation of the gluon density as a function of space and time in these approaches is very different from that in a simple Bjorken expansion. A previous calculation in this direction [7, 8] estimated the effects of 3-D expansion on the R_{AA} . However, this approach treated the energy loss of jets in a rather simplified manner, with the energy loss dE/dx exhibiting a simple linear dependence on the product of hard scattering cross section and gluon density (as a function of temperature extracted from the hydrodynamic simulation). Over the past year we have utilized our 3-D hydrodynamic model to provide the time-evolution of the medium produced at RHIC for jet energy-loss calculations performed in the BDMPS/ASW [9], Higher Twist [10] and AMY [11] approaches. In each of the three efforts, the inclusive as well as the azimuthally differential nuclear suppression factor R_{AA} of pions was studied as a function of their transverse momentum p_T . In addition, the influence of collective flow, variations in rapidity, and energy-loss in the hadronic phase were addressed for the selected approaches. For details regarding the implementation of the energy-loss schemes and their interface to the hydrodynamic medium, we refer the reader to the publications cited above. Here we shall focus on a systematic comparison between the three approaches, utilizing the same hydrodynamic medium evolution as well as the same structure and fragmentation functions for calculating the initial state and final high- p_T hadron distributions.

The left frame of figure 1 shows the nuclear modification factor R_{AA} in $Au-Au$

collisions at 0-5% (top) and 20-30% (bottom) centrality calculated in the ASW, HT and AMY approaches compared to data from PHENIX [12]. As can be seen, the parameters for all three approaches (initial maximal value for the transport coefficient $\hat{q}_0$ or coupling constant α_s in the AMY case) can be adjusted such that the approaches are able to describe the centrality dependence of the nuclear modification factor reasonably well. For a gluon jet, the values are $\hat{q}_0 \approx 3.4 \text{ GeV}^2/\text{fm}$ for the HT approach, $\hat{q}_0 \approx 20 \text{ GeV}^2/\text{fm}$ for the ASW formalism and $\alpha_s \approx 0.33$ for the AMY approach, which can be converted into a value of $\hat{q}_0 \approx 5.5 \text{ GeV}^2/\text{fm}$. Note that the ASW value for $\hat{q}_0$ at $\tau = 0.6 \text{ fm}/c$ and $\epsilon_0 = 55 \text{ GeV}/\text{fm}^3$ lies a factor of 3.6 higher than the Baier estimate for an ideal QGP, $\hat{q} \approx 2 \cdot \epsilon^{3/4}$ [13], while the AMY value is in line with the ideal QGP estimate and the HT calculation lies about a factor of 1.6 below that estimate. The large difference in $\hat{q}_0$ values between HT and ASW has been pointed out previously. However, we find that a factor of two can be accounted for by the use of different scaling prescriptions (temperature vs. energy-density) with which the medium is coupled to the transport coefficient – this will be discussed in greater detail in a forthcoming publication.

We find that slight variations appear between the approaches when R_{AA} is studied as a function of azimuthal angle. This can be seen in the right frame of figure 1 where R_{AA} is plotted as a function of azimuthal angle at $p_T = 10 \text{ GeV}/c$ (solid line) and $p_T = 15 \text{ GeV}/c$ (dashed line) for all three approaches in the 20-30% centrality bin. In order to quantify the difference between the three approaches we calculate the ratio of the out of plane R_{AA} over the in plane R_{AA} as a function of transverse momentum – this is shown in the left frame of figure 2. We find that AMY and HT exhibit the same peak to valley ratio, even though the absolute values for R_{AA} differ by approximately 10%. The ASW calculation systematically shows a stronger azimuthal dependence than the HT and AMY calculations - the cause of which will require a more detailed analysis to determine.

In order to investigate the spatial response of the jet energy-loss schemes to the medium the right frame of figure 2 shows the escape probability of a hadron with a transverse momentum between 6 and 8 GeV/c originating from a quenched jet moving in the positive x direction in the transverse plane as a function of its production vertex along the x -axis. Mathematically this quantity is defined as:

$$P(x) = \int dy T_{AB}(x, y) \cdot R_{AA}(x, y) / \int dx dy T_{AB}(x, y) \cdot R_{AA}(x, y) \quad (1)$$

It is remarkable how well the three different approaches agree with each other in this quantity. Since the same hard scattering probability was used as input in all three cases, the agreement in $P(x)$ really shows that all three approaches yield the same suppression factor as a function of production vertex of the hard probe, i.e. that they probe the density of the medium in the same way.

In summary, our comparison shows that under identical conditions (i.e. same medium evolution, same choice of parton distribution functions, scale etc.) all three jet energy-loss schemes yield very similar results. This finding is very encouraging since it indicates that the technical aspects of the formalisms are well under control. However,




Figure 2. Left: Ratio R_{AA} for out of plane vs. in plane emission as a function of p_T at $b=7.5$ fm impact parameter for all three approaches. Right: escape probability of a hadron with 6-8 GeV/c transverse momentum moving along the positive x -axis in the transverse plane as a function of x .

we need to point out that there still exist significant differences regarding the extracted value for the transport coefficient $\hat{q}_0$, which have yet to be fully understood.

Acknowledgments

This work was supported in part by an Outstanding Junior Investigator Award from the U. S. Department of Energy (grant DE-FG02-05ER41367). TR was supported in part by the Academy of Finland, Project 206024 and CG, GQ and JR acknowledge support by the Natural Sciences and Engineering Research Council of Canada.

References

- [1] Adcox K *et al.* (PHENIX) 2002 *Phys. Rev. Lett.* **88** 022301 (*Preprint nucl-ex/0109003*)
- [2] Adler C *et al.* (STAR) 2002 *Phys. Rev. Lett.* **89** 202301 (*Preprint nucl-ex/0206011*)
- [3] Gyulassy M and Wang X n 1994 *Nucl. Phys.* **B420** 583–614 (*Preprint nucl-th/9306003*)
- [4] Baier R, Dokshitzer Y L, Mueller A H, Peigne S and Schiff D 1997 *Nucl. Phys.* **B483** 291–320 (*Preprint hep-ph/9607355*)
- [5] Zakharov B G 1997 *JETP Lett.* **65** 615–620 (*Preprint hep-ph/9704255*)
- [6] Nonaka C and Bass S A 2007 *Phys. Rev.* **C75** 014902 (*Preprint nucl-th/0607018*)
- [7] Hirano T and Nara Y 2002 *Phys. Rev.* **C66** 041901 (*Preprint hep-ph/0208029*)
- [8] Hirano T and Nara Y 2004 *Phys. Rev.* **C69** 034908 (*Preprint nucl-th/0307015*)
- [9] Renk T, Ruppert J, Nonaka C and Bass S A 2007 *Phys. Rev.* **C75** 031902 (*Preprint nucl-th/0611027*)
- [10] Majumder A, Nonaka C and Bass S A 2007 *Phys. Rev.* **C76** 041902 (*Preprint nucl-th/0703019*)
- [11] Qin G Y *et al.* 2007 *Phys. Rev.* **C76** 064907 (*Preprint arXiv:0705.2575[hep-ph]*)
- [12] Shimomura M (PHENIX) 2006 *Nucl. Phys.* **A774** 457–460 (*Preprint nucl-ex/0510023*)
- [13] Baier R 2003 *Nucl. Phys.* **A715** 209–218 (*Preprint hep-ph/0209038*)